

Evidence supporting exercise-induced pulmonary haemorrhage in racing greyhounds

TS Epp^{1,*}, B Szladovits¹, A Buchannan¹, L Gates¹, P McDonough², DJ Padilla¹, J Smart³, HH Erickson¹ and DC Poole¹

¹Department of Anatomy and Physiology, Kansas State University, Manhattan, Kansas 66506, USA

²Department of Kinesiology, University of Texas at Arlington, Arlington, Texas 76019, USA

³Veterinary Consultant for the National Greyhound Racing Association, 619 Woodland Drive, Washington, Kansas 66968, USA

* Corresponding author: tepp@vet.k-state.edu

Submitted 23 May 2007; Accepted 8 January 2008

Research Paper

Abstract

Exercise-induced pulmonary haemorrhage (EIPH) is a major health concern in performance horses, but the presence and severity of this condition in racing greyhounds has received little attention. While equids and greyhounds share many physiological attributes, there are important structural and functional differences that may help protect greyhounds from EIPH. We tested the hypothesis that greyhounds performing a simulated 503 m race would experience EIPH and that the time course of recovery would be similar to the horse, even though the severity or relative extent as indexed by the concentration of red blood cells [RBCs] in bronchoalveolar lavage (BAL) fluid would be lower in comparison with that demonstrated previously in horses. Greyhound dogs ($n = 6$) raced on two occasions (separated by 7 weeks) and BAL was performed 1 week before, 2 h after and each week for 4 weeks following each race to examine the [RBC], concentration of white blood cells [WBCs], WBC differentials and haemosiderophages in the lungs. Racing increased 10 min post-exercise venous blood [lactate] to $18.6 \pm 0.4 \text{ mmol l}^{-1}$. No epistaxis or pink froth was observed at the nose or mouth of any of the dogs. The [RBC] in the BAL fluid was increased significantly 2 h post-race (baseline = $109.6 \pm 11.7 \times 10^3$; post-race = $292.3 \pm 69.9 \times 10^3 \text{ RBC ml}^{-1}$ BAL fluid, $P < 0.05$) and returned to baseline 1 week post-race ($149.2 \pm 46.2 \times 10^3 \text{ RBC ml}^{-1}$ BAL fluid, $P > 0.05$ versus baseline). The number of haemosiderophages was not different for any of the measurement periods. The [WBC] in the BAL fluid decreased from baseline and race values at 2, 3 and 4 weeks post-exercise (all $P < 0.05$). Alveolar neutrophil concentrations were also decreased from baseline and immediate post-race values for 4 weeks post-race. The increased [RBC] in the BAL fluid post-exercise is consistent with the presence of EIPH in these greyhounds. However, the relative extent of EIPH in greyhounds (as indexed by [RBC] in the BAL fluid), as compared with that in the horse, was mild, and the lack of elevation of WBC suggests that, unlike their equine counterparts, inflammatory airway disease was absent.

Keywords: EIPH; racing greyhounds; bronchoalveolar lavage; inflammatory airway disease

Introduction

Exercise-induced pulmonary haemorrhage (EIPH) has been reported to occur in nearly all horses¹⁻⁴, as well as camels⁵, humans^{6,7} and dogs^{8,9} following strenuous exercise. Although this condition has been well documented in the Thoroughbred horse since the 17th century¹⁰ and researched extensively, only two preliminary reports provide evidence that greyhounds experience EIPH^{8,9}.

In horses, the aetiology of EIPH is multifactorial, with high pulmonary artery pressures^{11,12}, large airway pressure swings^{3,13-22}, inflammatory airway disease (IAD)^{1,23-29} and ground impact forces³⁰⁻³² hypothesized to be the major contributing factors. The exercise response of greyhounds is physiologically similar to that of the horse in many ways. The Thoroughbred horse and racing greyhound are both exceptional sprinters³³⁻³⁵ (c. 18 m s^{-1}) with massive cardiac outputs³³⁻³⁶ (0.6 and $1.01 \text{ kg}^{-1} \text{ min}^{-1}$ in horse and greyhound, respectively)

and exceptional elevations in packed cell volume (PCV; 65 and 66% in horse and greyhound, respectively)^{34,35,37,38} during maximal exercise. The pulmonary capillary transmural pressure required for rupture in the canine pulmonary capillary (66–70 mmHg)^{39,40} measured *ex vivo* is similar to the mean pulmonary artery pressure (Ppa) reported for the greyhound during sub-maximal exercise at 11 m s⁻¹ (c. 55 mmHg)⁴¹. Thus, it is likely that the greyhound approaches or exceeds this ‘threshold for capillary rupture’ during maximal exertion. One important difference between canine and equine athletes is that horses are obligate nasal breathers during exercise. This, in combination with extraordinarily high airflow rates and a long trachea (where length of airway is proportional to resistance of airway according to Poiseuille’s law), causes the development of exceptionally sub-atmospheric extravascular pulmonary pressures, which summate with the intravascular pressures to create the very high transmural pressures measured in the horse. As the greyhound is not an obligate nasal breather and during exercise achieves far lower peak airflows than the horse through a much shorter, though narrower, trachea, it is possible that the fall in alveolar pressure^{42–44} is less extreme and this therefore reduces the potential for EIPH in this species.

While pulmonary capillary transmural pressures may be lower in greyhounds than in horses, the pulmonary capillary blood–gas barrier is weaker in the dog⁴⁰. Specifically, when subtracting oesophageal pressure (Pes) from mean Ppa to estimate pulmonary artery transmural pressure, a value of greater than 65 mmHg (close to canine stress failure threshold) is estimated during sub-maximal exercise. Thus, it should not be surprising that preliminary data suggest that greyhounds do experience EIPH during maximal sprint exercise^{8,9}.

This investigation was designed to determine whether greyhounds run over the standard 503 m course demonstrate a significant increase in [RBC] in the BAL fluid, and, if present, outline the timeline for resolution. Specifically, we tested the hypothesis that greyhounds would exhibit a significant increase in [RBC] in the BAL fluid, though to a lesser relative extent in comparison with the horse, but following a similar pattern and time course of recovery (i.e. disappearance of [RBC], elevation of white blood cells (WBC) and haemosiderophage emergence, peak and decline) to that documented for the horse.

Methods

Animals

Six healthy greyhound dogs that had been raced on the track were acquired for this study. The group consisted of four intact females and two intact males ranging in

age from 2 to 4 years and weighing 29.0 ± 1.2 kg. They were housed in a temperature-controlled building (70–74°F with 30–70% humidity). They had a 0.3 m² area inside their pen with free access to a 74 m² outdoor sand run. Indoor conditions were on a 12-h light:12-h dark cycle. The dogs were fed Iams (Iams Mini Chunks®, Iams Company, Dayton, OH, USA) adult dog food (minimum of 26% crude protein, minimum of 15% crude fat, maximum of 5% crude fibre and maximum of 10% moisture) once daily in the morning and had free access to water. The greyhounds were current on vaccinations including distemper virus, adenovirus, parainfluenza virus, parvovirus and *Bordetella bronchiseptica*, as well as being on monthly Heartgard Plus. All procedures used were approved by the Kansas State University Animal Care and Use Committee.

Animal preparation

The greyhounds were fed 3 h before estimated race time. The dogs were transported in a climate-controlled dog trailer c. 81 km to a local training track.

Exercise protocol

Greyhounds were divided into two heats with three dogs in each heat and run on the track in a race-simulated timed race (503 m distance) around a track, chasing a lure. After 9–10 min of completing the race, a venous blood sample was obtained from the jugular vein with a 22 gauge 1 in. needle attached to a 3 cc heparinized syringe with a stopcock for post-exercise plasma lactate analysis. Greyhounds were not formally exercised post-race, but did have access to their runs. A second identical race (race 2) was performed 2–3 weeks after RBC and WBC counts had stabilized near baseline levels (c. 7 weeks post-race 1).

Bronchoalveolar lavage

The initial baseline BAL was performed and cultures obtained after the dogs had not been formally exercised or raced for 1–2 weeks after their acquisition by Kansas State University. One week after the baseline lavage, the initial post-race lavage was performed c. 2 h post-exercise under general anaesthesia, this sampling procedure examined the number of RBC per millilitre of BAL fluid (which is considered to be a reflection of the amount of blood in the airways post-exercise), which we took as an indirect measure of the relative extent of EIPH^{2,3,8,9,17–19,22,23,45–49}. Briefly, after intramuscular pre-medication with atropine (0.025 mg kg⁻¹), morphine (0.25 mg kg⁻¹) and diazepam (0.2 mg kg⁻¹) had taken effect, the dogs were induced with intravenous propofol (6 mg kg⁻¹; Propoflo™, Abbott Laboratories, North Chicago, IL, USA), intubated and placed on supplemental oxygen. Dogs remained in sternal recumbency for the

procedure. They were maintained with an intravenous infusion of propofol ($0.4 \text{ mg kg}^{-1} \text{ min}^{-1}$) in order to allow the collection of the BAL fluid samples. For health and safety reasons, heart rate, systemic arterial pressure, oxygen saturation, respiratory rate/character, mucous membrane colour and depth of anaesthesia were monitored by a technician. After the animals were stabilized under anaesthesia, a flexible modified foal stomach tube⁵⁰ with a diameter of 16-F (Argyle stomach tube, Sherwood Medical, Company, St Louis, MO, USA) was wedged in a sub-segmental bronchus of the caudal dorsal lung lobe after insertion through the endotracheal tube. The location of the distal end of the BAL tube was assumed to be the dorsocaudal lung region from the work of Hawkins⁵⁰, but it was verified radiographically on all six greyhounds before and after one of the lavage trials. Before insertion, the distal end was transected with a sterile blade above the most proximal fenestration (c. 6 cm from the distal end). A sterile hand-held metal pencil sharpener was used to round the distal end of the tube and create a slight taper to enhance formation of a tight seal when the tube was wedged in a bronchus of the caudal lung lobe. During insertion once resistance was felt, an attempt was made to gently withdraw and reposition the catheter, ensuring placement in the most distal airway possible. A Christmas tree adapter was placed on the proximal end of the BAL catheter. Following placement of the catheter, a 50 ml aliquot of sterile isotonic saline (0.9%), warmed to body temperature, was slowly infused. After a short delay (c. 2 breaths), the fluid was aspirated with gentle suction while chest copage was performed bilaterally. This procedure was repeated thrice, equivalent to a dose of 5 ml kg^{-1} of instilled fluid per dog (c. 125–150 ml of total per dog). Supplemental oxygen was administered after the procedure until extubation and thereafter provided by mask until the dog recovered. The dogs were monitored every 4–6 h after the procedure for signs of respiratory distress, pale mucous membranes or prolonged capillary refill time, and twice daily for the next 48 h for signs of respiratory distress, cough, lethargy or loss of appetite. The BAL samples were placed on ice immediately after collection and remained chilled throughout the analysis. The BAL fluid was centrifuged (10 min at $600 \times g$; TJ-6 Table Top Centrifuge, Beckman Instruments, Inc., Palo Alto, CA, USA), after which the supernatant was removed via gentle suction and the pellet resuspended in sterile 0.9% saline. RBC and WBC were then quantified manually with standard haemocytometer counts (Fisher Scientific, Pittsburgh, PA, USA; Nikon, Inc., Instrument Group, Garden City, NY, USA), expressed as cells ml^{-1} BAL fluid, and differential slides were made (Cytospin 2, Shandon, Pittsburgh, PA, USA) and evaluated. These slides

were stained with a diff-quick stain (Hema-Stain, Fisher Scientific) that stains cells similar to the Wright–Giemsa stain. Cytospin slides were also stained with Perl's Prussian Blue Stain as previously described by Meyer *et al.*² and the number of haemosiderophages was determined from the entire slide that had uniform number of cells (1×10^6) per slide.

After the initial post-race lavage, the lavage procedure was then performed at 1-week intervals to determine the time course of recovery from EIPH. There was a 2-week interval between the fourth-week post-race lavage and baseline 2. The post-race 2 lavage was done 1 week after the second baseline in a similar manner to race 1.

Blood analysis

Following venous blood withdrawal (jugular vein c. 10 min post-race) into plastic heparinized syringes, blood samples were placed immediately on ice. Within 2 h of the experiment, plasma lactate was quantified using a Nova Stat M blood-gas analyzer (Nova Biomedical, Waltham, MA, USA). This machine was calibrated according to the manufacturer's standards immediately prior to running the samples.

Statistics

Differences in measured variables were analysed using a mixed effects model where time is a fixed effect, and dog and dog \times time are random effects. When significant differences were found, a least-square means post-hoc test was used to determine where the differences existed. The Statistical Analysis System Program 9.1.2 statistical package (SAS Institute, Inc., Cary, NC, USA) was used to analyse the data. Significance was accepted at the $P \leq 0.05$ level.

Results

The percentage of BAL fluid recovery was similar for all lavages (race and resting lavages) and ranged from 59 to 68% with an average of $63 \pm 1\%$ (Table 1). Cultures obtained from the first lavage were negative for any pathogens.

Red blood cell concentrations in the lavage fluid

The greyhounds exhibited a significant increase in the [RBC] in the BAL fluid ($P = 0.025$) when determined after both race 1 ($P = 0.036$) and race 2 ($P = 0.016$). Baseline and post-race 1 values (mean \pm SE) were $144.5 \pm 21.9 \times 10^3$ and $313.8 \pm 70.5 \times 10^3$ RBC ml^{-1} BAL fluid, respectively. Baseline and post-race 2 values (mean \pm SE) were $74.7 \pm 13.1 \times 10^3$ and $270.7 \pm 120.1 \times 10^3$ RBC ml^{-1} BAL fluid, respectively. The mean levels of RBC ml^{-1} in the BAL fluid for the baselines and races were consistent (no significant differences existed between baseline 1 *versus* baseline

Table 1 Bronchoalveolar lavage (BAL) data

Variable	Baseline	Race	1 WPR	2 WPR	3 WPR	4 WPR
Total RBC ml ⁻¹ BALF	109 600 ± 11 700	292 300 ± 69 900*	149 200 ± 46 200†	79 000 ± 15 500†	94 900 ± 31 100†	109 900 ± 57 700†
Total WBC ml ⁻¹ BALF	134 000 ± 14 900	141 000 ± 19 900	122 000 ± 32 100	84 000 ± 10 900†*	89 700 ± 14 500†*	75 200 ± 39 000†*
% Return BALF	66.3 ± 3.5	60.7 ± 3.1	63.6 ± 4.4	59.5 ± 3.4	60.3 ± 6.0	66.7 ± 4.3
Macrophages (cells ml ⁻¹ BALF)	93 100 ± 9330 (70.7 ± 3.3)	95 600 ± 14 200 (68.6 ± 2.9)	95 500 ± 4960 (78.3 ± 4.1)*†	64 600 ± 2090 (76.5 ± 2.5)†	66 900 ± 3100 (74.5 ± 3.5)	50 000 ± 3960 (66.5 ± 5.3)
Neutrophils (cells ml ⁻¹ BALF)	17 000 ± 3250 (12.6 ± 2.5)	16 400 ± 4020 (11.3 ± 2.1)	7310 ± 2670*† (6.0 ± 2.2)	7740 ± 1470*† (9.2 ± 1.7)	7920 ± 1260*†(8.8 ± 1.4)	9770 ± 2990*†(13.0 ± 4.0)
Lymphocytes (cells ml ⁻¹ BALF)	19 900 ± 3840 (14.3 ± 1.7)	26 600 ± 5080 (18.7 ± 2.4)	18 700 ± 3600 (15.3 ± 3.0)	11 800 ± 2040 (14.0 ± 2.4)	14 810 ± 2240 (16.5 ± 2.5)	13 800 ± 2220 (18.3 ± 3.0)
Eosinophils (cells ml ⁻¹ BALF)	44 10 ± 2920 (2.4 ± 1.7)	22 50 ± 1660 (1.6 ± 1.0)	610 ± 610 (0.5 ± 0.5)	279 ± 279 (0.3 ± 0.3)	153 ± 153 (0.2 ± 0.2)	1630 ± 1630 (2.2 ± 2.2)
Mast Cells (cells ml ⁻¹ BALF)	214 ± 163 (0.2 ± 0.1)	0 ± 0	0 ± 0	144 ± 144 (0.2 ± 0.2)	0 ± 0 (0 ± 0)	0 ± 0 (0 ± 0)
Haemosiderophages	41 ± 10	64 ± 38	108 ± 40	80 ± 36	21 ± 4	77 ± 38

Data represent averaged bronchoalveolar lavage (BAL) variables for six greyhounds that were lavaged 2 h, and weekly, after two separate 5/16 mile races. The differential counts are presented with the actual number of white blood cells (WBCs) by type (i.e. macrophage, neutrophils, lymphocytes, eosinophils and mast cells) on the top and the differential percentage directly below it. Haemosiderophage numbers represent the number on a slide with 1×10^6 WBCs per slide. Time periods across the top of the table are abbreviated as follows: baseline, resting BAL; 1 WPR, 1 week post-race BAL (resting); 2 WPR, 2 weeks post-race BAL (resting); 3 WPR, 3 weeks post-race BAL (resting); 4 WPR, 4 weeks post-race BAL (resting). *Significant difference ($P < 0.05$) from baseline; †Significant difference ($P < 0.05$) from race.

2 or race 1 *versus* race 2), so these two data points were averaged (baseline = $109.6 \pm 11.7 \times 10^3$ RBC ml⁻¹ BAL fluid and race = $292.2 \pm 69.9 \times 10^3$ RBC ml⁻¹ BAL fluid) for ease of presentation and interpretation (Fig. 1). The number of RBC ml⁻¹ BAL fluid had returned to baseline by 1 week post-race (149.2 ± 46.2 RBC ml⁻¹ BAL fluid; Fig. 1).

Additional bronchoalveolar lavage fluid analysis

As there was no significant difference between the two races, the baseline and race lavages were averaged as for the RBC data. There was no change in the [WBC] 1 week post-race, but there was a significant decrease from baseline 2–4 weeks post-race (Fig. 2; Table 1). Haemosiderophage counts for these periods were quite variable but did not change significantly from baseline (Table 1). Despite the absence of an inflammatory response, there were significant differences in the differential counts. Specifically, the concentration of neutrophils in the BAL fluid was decreased from baseline during the entire 4 weeks post-race ($P < 0.05$, Table 1). As a result of these decreased neutrophils, the relative percentage of alveolar macrophages (occasionally containing vacuolation) increased at 1 week post-race and remained significantly elevated for 2 weeks, returning to baseline levels at 4 weeks post-race (Table 1).

Blood lactate, rectal temperature and performance times

Venous blood [lactates] were 19.2 ± 0.4 and 18.0 ± 0.5 mmol l⁻¹ ($P > 0.05$) for 10 min post-race samples for races 1 and 2, respectively, with the average being 18.6 ± 0.4 mmol l⁻¹. Ten minutes post-race, rectal temperatures were 106.0 ± 0.3 and $104.5 \pm 0.4^\circ\text{C}$ for races 1 and 2, respectively ($P > 0.05$), with the average being $105.2 \pm 0.3^\circ\text{C}$. Race times were 32.1 ± 0.3 and 32.6 ± 0.4 s for races 1 and 2, respectively ($P > 0.05$, average 32.2 ± 0.1 s; a competitive time for the lower-end greyhound would be 31.0–31.5 s).

Discussion

The principal original findings of this investigation are that: (1) These greyhound dogs run under simulated race conditions at 503 m experienced an increase in [RBC] in post-race BAL, and that increase was of a lower magnitude compared with that described previously in horses^{2,17–20}. (2) Recovery from elevated [RBC] in the BAL fluid post-race in greyhounds followed a similar pattern to horses when looking at [RBC], in that the numbers of RBC ml⁻¹ BAL fluid returned to baseline levels within 1 week post-race. In contrast to the horse, this did not result




Fig. 1 Exercise-induced pulmonary haemorrhage (EIPH) in greyhounds as assessed by bronchoalveolar lavage (BAL; $n = 6$) performed before and after a 5/16 mile race. EIPH is expressed as the number of red blood cells (RBCs) per millilitre of BAL fluid. Data are presented as mean \pm SE of the average of the baselines (1 and 2) and the races (1 and 2). Time periods on the abscissa are abbreviated as follows: baseline, resting BAL; race, 2 h post-race BAL; 1 WPR, 1 week post-race BAL (resting); 2 WPR, 2 weeks post-race BAL (resting); 3 WPR, 3 weeks post-race BAL (resting); 4 WPR, 4 weeks post-race BAL (resting). Lavages were performed 1 week apart. *Significant increase in EIPH (increased number of RBC) from baseline; †significant decrease in the number of RBC per millilitre of BAL fluid compared with immediately post-race values

in prolonged elevation of haemosiderophages or an extended inflammatory response in the lung.

Comparison with the current literature

A very limited amount of information exists concerning EIPH in racing greyhounds (see preliminary reports)^{8,9}. King^{8,9} reported significant EIPH from the dorsocaudal lung region in greyhounds racing 503 m, ranging from 1 to 71 million RBC ml⁻¹ BALF with an occasional dog displaying endoscopic evidence of EIPH. The data from the current study are in agreement with the preliminary studies by King *et al.*^{8,9}, suggesting that greyhounds may demonstrate pulmonary haemorrhage during exercise. However, the [RBC]

detected in the BAL fluid in the current study was much lower than that found by King^{8,9}. Between-study differences in the [RBC] of the BAL fluid post-race may be due to methodological factors including time from run to lavage (20 min in King *et al.*^{8,9} studies *versus* 2 h in present study), induction of iatrogenic haemorrhage, lung differences (right *versus* left lung predominance in greyhounds) and pre-selection of dogs for the study (certain populations of animals may potentially be more prone to severe EIPH).

The greyhounds in the current study demonstrated an increase in [RBC] in the BAL fluid detected 2 h post-race, which returned to baseline levels within 1 week post-exercise as reported for the horse².




Fig. 2 White blood cells (WBCs) per millilitre of BAL fluid measured in the bronchoalveolar lavage (BAL) fluid 2 h after greyhounds ran a 5/16 mile race and at weekly intervals thereafter (abscissa labelled as for Fig. 1). Data are presented as mean \pm SE of the average of the baselines (1 and 2) and the races (1 and 2). Lavages were performed 1 week apart. No significant differences occurred from baseline to the race. *Significant decrease in WBCs per millilitre of BAL fluid post-race; †significant decrease in the number of WBCs per millilitre of BAL fluid from baseline ($P < 0.05$)

However, the amount detected in the airways is far lower than values measured typically in the horse (EIPH = $4\text{--}64 \times 10^6$ RBC ml⁻¹ BAL fluid)^{19,22,23,45-47,50}. Haemosiderophages were also examined in the study of Meyer *et al.*² as an indicator of past pulmonary haemorrhage^{1,2,51,52}. In horses, haemosiderophages are commonly around 7% at baseline and increase significantly (10-20%) 1 week post-exercise, remaining elevated for 3-4 weeks post-exercise. This equine profile is in contrast to the extremely low, almost undetectable levels of haemosiderophages that we found in greyhound dogs. Due to extremely sparse numbers of haemosiderin-laden macrophages, it was impossible to obtain a percentage of haemosiderophage count for the dog slides. Therefore, the total number of haemosiderophages per million cells (entire slide) was counted to increase the accuracy of comparison within dogs and among BAL periods. The small amount of blood detected in our study did not induce an increase in haemosiderophages 1-3 weeks post-exercise as observed previously in the horse². Moreover, alterations in the haemosiderophage counts for the greyhounds were erratic with respect to pre- and post-race evaluations, and this profile may potentially be explained by mild iatrogenic bleeding initiated during the lavage procedure. These data are in agreement with King *et al.*'s⁹ observation that 7 out of 10 dogs showed evidence of haemosiderin deposits within alveolar macrophages indicating past haemorrhage.

IAD in the horse is hypothesized to contribute to the initiation and severity of EIPH as well as being a response to the condition^{1,24-28}. A long-standing inflammatory reaction to blood within the alveoli results in a self-perpetuating phenomenon of gradually worsening haemorrhage as horses continue to train and race^{24,53}. However, in the racing greyhound, a much lower [RBC] in BAL was detected after a race, which may not be sufficient to overwhelm pulmonary defences and these RBCs may be cleared too rapidly to invoke any prolonged inflammatory response. It has been shown in the horse that alveolar macrophage function (phagocytosis and oxidative burst) becomes suppressed for 2-3 days when challenged by alveolar haemorrhage^{24,54-60} as a consequence of the limited ability of alveolar macrophages to effectively metabolize/detoxify the iron in RBC^{61,62}. It is also possible that IAD itself may not induce the long-term pulmonary damage that exacerbates EIPH in the horse^{27,29}. The present investigation did not find an elevated WBC concentration in the lavage fluid at baseline (in comparison with the literature)⁶³⁻⁷² nor following racing. However, an unexpected decrease occurred in the WBC concentration at weeks 2, 3 and 4 after racing. A decrease in the concentration of neutrophils was detected for 4 weeks post-race, which follows the

WBC concentration and corroborates the absence of an inflammatory response to the mild haemorrhage. It is possible that weekly use of propofol (an anaesthetic drug with anti-inflammatory properties) could aid in masking a mild inflammatory response^{73,74}. We did note a significant rise in the percentage of macrophages (not actual numbers) from 1 to 3 weeks post-race that resulted from a significant decrease in neutrophils, since the actual concentration of macrophages remained unchanged. These results are also in agreement with EIPH studies of King *et al.*^{8,9} but at variance with the equine literature where the WBC concentration typically becomes elevated after an episode of EIPH and either remains elevated or decreases slowly, depending on the independent contribution from IAD. Again, the greyhound dogs in the present investigation did not appear to have true IAD at any time (i.e. elevation in WBC concentration with concurrent increase in macrophages, neutrophils and/or lymphocytes). Other conditions cause inflammation in the lower airways of dogs, but are totally unrelated to the current study and include 'ski asthma' in Alaskan sled dogs, allergic bronchitis and eosinophilic airway disease⁷⁵. When looking at WBC concentration and differentials from BAL fluid, much variation exists in the canine literature. In support of our conclusion that a post-race inflammatory response was not evident, the WBC concentration and differentials obtained herein fall within the published ranges⁶³⁻⁷².

Mechanisms of exercise-induced pulmonary haemorrhage

EIPH has been studied most extensively in the Thoroughbred horse and several mechanisms have been proposed to contribute to the pathophysiology of EIPH in that species. In order of potential importance as considered in the literature, these include elevated pulmonary arterial pressures^{11,12,76}, airway contributions^{3,13-22}, IAD^{1,23-29}, blood viscosity⁷⁷ and locomotory impact forces³⁰⁻³².

Greyhounds may be expected to demonstrate EIPH as they have a number of physiologic attributes in common with the horse. Both species are exceptional sprinters^{33,34}, have large heart-to-body weight ratios⁷⁸⁻⁸¹, large contractile spleens (high PCV)^{34,37,38,77}, high pulmonary vascular pressures during exercise^{12,41} and very high maximal oxygen consumptions^{33,35,36,82,83} for their body size. In addition, greyhounds have a thinner blood-gas barrier than the horse that requires a smaller transmural pressure gradient to cause capillary rupture^{3,39,40,84}, as reported by Birks *et al.*⁸⁴ (dog, 0.795 ± 0.084 μm ; 66 mmHg and horse, 0.930 ± 0.044 μm ; 70-100 mmHg) or potentially similar when reported as harmonic mean thickness (dog $0.509 \pm 0.002 \times 10^{-4}$ cm and pony $0.511 \pm 0.031 \times 10^{-4}$ cm) by Weibel *et al.*⁸⁵. Greyhounds have high cardiac outputs^{35,34,36} that play a large role

(in combination with their highly elevated PCV) in generating both exceptional levels of oxygen uptake^{33,36} and potentially high pulmonary arterial pressures during maximal exercise. Based on pulmonary artery pressures of greyhounds measured during sub-maximal exercise (55 mmHg at 11 ms⁻¹), peak exercising pressures at speeds approaching 18 ms⁻¹ might well be much higher.

Notwithstanding the above, these are features specific to the greyhound that may protect this animal from the extensive rupture of the blood-gas barrier observed in the horse. Specifically, the dog is not an obligate nasal breather and has a much shorter trachea than the horse (airway resistance is proportional to the length of the airway and inversely proportional to the radius to the fourth power), thus, depending on the airflows achieved and airway(s) resistance, the work of breathing and pleural pressure swings generated⁴² may not be as high as those seen in the horse. Saibene *et al.*⁴³ found that dogs during exercise experienced a pronounced dilation of the respiratory tract, which reduces airflow resistance. The horse also has a very stiff chest wall that limits the thoracic displacement to ventilation during exercise⁸⁶ and mandates a proportionally greater diaphragmatic displacement. Consequently, horses may be predisposed to more sub-atmospheric intrapulmonary pressures and also shear forces that damage the dorsocaudal lung lobes. The more compliant chest wall of the dog may well limit these effects if it permits lateral chest wall expansion during exercise. Findings support the hypothesis that impact forces are a significant risk factor for EIPH in horses³⁰⁻³². However, these forces will be reduced in the dog in proportion to their body frames and weights, along with the consideration that they are not carrying a rider and saddle. Lastly, the severity of EIPH may depend on the duration over which the threshold capillary transmural pressure is exceeded⁸⁷. Thus, the shorter distance (greyhounds, 503 m *versus* Thoroughbred horses, 1609–2416 m) and reduced time (25–35 s for greyhounds *versus* 1–2 min for horses) at which maximal effort is sustained by the greyhound *versus* the Thoroughbred horse may help constrain the magnitude of EIPH. However, racing Quarter Horses can demonstrate extreme EIPH in races of similar distance to greyhounds⁸⁸, and Thoroughbreds themselves can evidence epistaxis breaking from the starting gate.

Methodological considerations

Bronchoalveolar lavage (BAL) has been utilized for a little over a decade in horses to evaluate the concentration of RBC present in the BAL fluid post-exercise⁵¹, and is frequently referred to in the equine^{2,3,17-19,22,45-49} and human literature^{7,89} to quantify the degree or

severity of EIPH. An advantage of BAL is that there is not a precise window of time within which BAL has to be performed, as is the case with endoscopy^{2,90}. In addition, there is less chance of missing blood due to the timing of post-exercise endoscopy or failure of blood movement into the larger airways where it can be visualized and graded^{2,48,91}. Even though King *et al.*^{8,9} did visualize small amounts of bleeding in two dogs with endoscopy, the relatively low [RBC] in BAL observed post-race in the current study would not likely have been observed with endoscopy^{1,2,51,92}. It is possible that these low [RBC] found in these greyhounds may be partially explained by the 75- to 90-min delay in performing the lavage procedure. The 2 h post-race lavage was chosen *versus* 30 min post-race as is conventional in the equine literature^{2,3,17-19,22,45-48} and similar to the King studies^{8,9} due to the increased safety of transporting the dogs back to the veterinary teaching hospital for the procedure instead of anaesthetizing the dogs at the track. It has been demonstrated that smaller RBC counts may be observed with large time lapses post-race, as in the study of Valdez *et al.*⁴⁹ where BAL was performed 12–18 h post-race. On the other hand, the level of bleeding could truly be less in maximally exercised dogs and may be further lowered by the extended time between race and lavage.

BAL has been considered to be the most accurate^{2,51} and sensitive⁹¹ technique available to detect and quantify the level of pulmonary haemorrhage. However, an obvious limitation of equating [RBC] with severity is its reliability, because this assumes that RBC counts in the lavage fluid are proportional to the severity of haemorrhage, which has not been definitively shown⁹¹. Despite this inability to absolutely quantify the amount of blood in the lung, utilization of the BAL technique and enumeration of [RBC] offers the best estimate of the extent of EIPH aside from removing the entire lung for whole lung lavage¹⁹, which is impossible for repeatedly quantifying the severity of EIPH in living horses⁹¹. Furthermore, techniques such as scintigraphy and endoscopy have greater limitations to accuracy and sensitivity and limit quantitative and objective assessment¹⁹. What has been shown, at least in horses where EIPH has been studied extensively, is that a good correlation exists between BAL cytology and histopathology in horses with EIPH^{49,92,93}.

It also has been shown in the horse that blind placement of the BAL tube typically results in wedging it in the dorsocaudal region⁹⁴ of the lung where the majority of EIPH occurs^{25,26,51,94}. Due to physiological similarities between the dog and the horse, as well as the preliminary data gathered with the bronchoscope in the greyhound^{8,9}, the dorsocaudal region was the site the authors chose to examine, realizing that the

differences in canine physiology and anatomy could potentiate bleeding in other locations of the canine lung. Unpublished radiographic data (TS Epp, AM Buchannan, L Gates, DC Poole and HH Erickson) from our laboratory suggest a similar caudal lung lobe placement (seven out of seven dogs) when a BAL tube is passed blindly (five right caudal lung lobes *versus* two left caudal lung lobes) in the greyhound. Even though side predominance is not statistically significant, this is in agreement with Hawkins *et al.*⁵⁰, who radiographically demonstrated right caudal lung lobe predominance when a BAL tube was blindly placed (seven dogs right *versus* two dogs left). Therefore, it may be of importance that King *et al.*^{8,9} showed that the left lung may haemorrhage to a greater degree than the right one, which is in contrast to what has been observed in the horse^{2,95}. It is also important to keep in mind that the greyhounds utilized in this study were not selected based upon a prior history of severe EIPH, as is often the case in horse studies^{3,19,22,23,45-47}. In addition, though these greyhounds raced giving maximal effort, their racing times were 0.7-1.0s slower than many competitive greyhounds, which may have impacted the relative extent of EIPH.

BAL is becoming more popular as a diagnostic tool for pulmonary disorders in dogs. However, BAL results can be difficult to interpret. Disadvantages or problems potentially encountered in dogs (but not in horses) include the possibility that alveoli are non-uniform and intermittent collapse occurs, not allowing lavage fluid into the distal airways as well as an increased rate of mucociliary clearance in the dog *versus* the horse (3.3 *versus* 2 cm min⁻¹)⁹⁶⁻⁹⁸. In addition, there may be mild trauma to the airways obtained while trying to ensure wedging of the tube^{50,67,68} and there is the potential for airway collapse⁶⁸ distal to the catheter if too much suction is inadvertently applied, resulting in possible trauma and a smaller recovery volume. In the current study, precautions were taken to minimize this trauma (i.e. use of sufficiently rigid tubing so that resistance could be felt and to prevent bending upon itself, but flexible enough to pass through the airways and minimize trauma with careful passage; continuous gentle suction applied with cessation upon first indication of negative pressure during aspiration of sample). It is not possible to differentiate whether RBC detected in the lung lavages originated from EIPH or from iatrogenic bleeding, but [RBC] are consistent across samplings from horses at rest². In the current study, baseline 1, weeks 1, 2, 3 and 4 post-race, as well as baseline 2 were performed 1 week apart with consistent [RBC], which would be unexpected if the techniques were inducing iatrogenic haemorrhage. There is no qualitative colour evidence of haemorrhage, but

baseline [RBCs] are present in horses not exercised (at least 10 days to 2 weeks)^{2,48,99}, so it is not surprising that baseline levels are also evident in the canine. The [RBC] in the BAL fluid has been used in horses as an index to indicate the relative extent of EIPH with various therapeutic interventions and alterations in exercise intensity and workload. The results and conclusions have been consistent, predictable and reproducible in multiple investigations from multiple laboratories. Lower [RBCs] are observed during sub-maximal (trotting) exercise *versus* maximal effort⁴⁸, at different exercise intensities³, and during inclined *versus* flat running⁴⁶. To date, BAL has demonstrated a reduction in EIPH with nasal strips^{17-19,22,49}, furosemide^{18,19,22}, and concentrated equine serum²³. An increased [RBC] in the BAL fluid has been demonstrated with nitric oxide inhibition using L-Name¹⁰⁰ and inhaled nitric oxide⁴⁵. Therefore, it follows from the equine data that the significant increase above baseline, which was observed in the [RBC] 2 h post-exercise in the dog, would be consistent with the occurrence of EIPH.

Considerable variation and lack of standardization exist in the precise volume of lavage fluid infused, the technique employed (i.e. the number of aliquots, interval between instillation and aspiration, lobes lavaged)^{71,101} and the processing of fluid (i.e. centrifugation, washing and counting)^{63,65,67,71}. The techniques utilized in this study were chosen after careful perusal of the available literature as well as some trial and error, and were found to combine the most consistent results with the least amount of iatrogenic trauma possible. Finally, depending specifically on volume and the number of aliquots, normal numbers and differential percentage of WBC in the lavage fluid are less defined than those available for equids¹⁰². Variations in the WBC counts as well as the differential cell counts^{65,71} may possibly be due to lavaging of different levels (i.e. bronchial *versus* alveolar)⁷², different concentrations obtained^{50,63} and the lung lavaged (i.e. left or right)⁸.

Implications of increased [RBC] in the BAL fluid of greyhounds post-exercise

The current investigation provides evidence that the greyhound exhibits a significant increase in [RBC] in the BAL fluid following maximal running, which is consistent with EIPH, albeit at very low levels compared with equids. However, one must use caution when directly comparing the [RBC] in the BAL fluid from equids to canids because any differences in pulmonary anatomy/physiology and the relative size of the scope or BAL tube to the airway could alter the level of haemorrhage detected in response to exercise. On the other hand, if these greyhounds are typical of

the entire racing population, it does not appear that the consequences of that bleeding are as pernicious and clinically worrisome as is the case for the horse. This conclusion is based on the fact that the levels of WBC, haemosiderophages and differential counts did not indicate a prolonged response to the blood (i.e. large inflammatory response and elevations in haemosiderophages over time) as is typically seen with the horse. Therefore, these dogs may not suffer the long-term lung damage and performance deficits created by inflammation and fibrosis. However, validation of this hypothesis will necessitate post-mortem pulmonary anatomical studies similar to those done in horses. Such studies are necessary to confirm or refute the notion that EIPH is not of major clinical significance in the greyhound as it is in the horse.

Acknowledgements

We are grateful to the following individuals who contributed significantly to this work: Dr Jim Higgins (statistics), Mr Maurie Flynn, Dr Bill Fortney, Dani Goodband, Susan Rose, Linda Schoen, Sandy Gammill, Marc Epp, Lindsey McClintock, Lee Buchannan, Nicole Harper, Katie Edwards and Rachel Lindberg.

This work received funding support from the Kansas Racing and Gaming Commission.

References

- 1 McKane SA, Canfield PJ and Rose RJ (1993). Equine bronchoalveolar lavage cytology: survey of Thoroughbred racehorses in training. *Australian Veterinary Journal* **70**: 401-404.
- 2 Meyer TS, Fedde MR, Gaughan EM, Langsetmo I and Erickson HH (1998). Quantification of exercise-induced pulmonary haemorrhage with bronchoalveolar lavage. *Equine Veterinary Journal* **30**: 284-288.
- 3 Langsetmo I, Fedde MR, Meyer TS and Erickson HH (2000). Relationship of pulmonary arterial pressure to pulmonary hemorrhage in exercising horses. *Equine Veterinary Journal* **32**: 379-384.
- 4 Birks EK, Shuler KM, Soma LR, Martin BB, Marconato L, Del Piero F, Teleis DC, Schar D, Hessinger AE and Uboh CE (2002). EIPH: posttrace endoscopic evaluation of Standardbreds and Thoroughbreds. *Equine Veterinary Journal Supplement* **34**: 375-378.
- 5 Akbar SJ, Derksen FJ, Billah AM and Werney U (1994). Exercise induced pulmonary hemorrhage in racing camels. *Veterinary Record* **135**: 624-625.
- 6 Hopkins SR, Schoene RB, Henderson WR, Spragg RG, Martin TR and West JB (1997). Intense exercise impairs the integrity of the pulmonary blood-gas barrier in elite athletes. *Journal of Applied Physiology* **79**: 908-917.
- 7 Hopkins SR (2005). The lung at maximal exercise: insights from comparative physiology. *Clinics in Chest Medicine* **26**: 459-468.
- 8 King RR, Raskin RE and Rosbalt JP (1990). Exercise-induced pulmonary hemorrhage in the racing greyhound dog. *Journal of Veterinary Internal Medicine* **4**: 130.
- 9 King RR and Raskin RE (1991). Intrabronchial hemorrhage in Greyhound dogs after racing. *7th International Racing Greyhound Symposium* Orlando, FL, in conjunction with Eastern States Veterinary Conference, 18-21.
- 10 Robertson JB (1913). Biological search-light on racehorse breeding: VI. The heredity of blood-vessel breaking in the Thoroughbred. *The Bloodstock Breeders Review* **2**: 265-281.
- 11 Erickson BK, Erickson HH and Coffman JR (1990). Pulmonary artery, aortic, and oesophageal pressure changes during high intensity treadmill exercise in the horse: a possible relation to exercise-induced pulmonary haemorrhage. *Equine Veterinary Journal Supplement* **9**: 47-52.
- 12 West JB, Mathieu-Costello O, Jones JH, Birks EK, Logemann RB, Pascoe JR and Tyler WS (1993). Stress failure of pulmonary capillaries in racehorses with exercise-induced pulmonary hemorrhage. *Journal of Applied Physiology* **75**: 1097-1109.
- 13 Jackson JA, Ducharme NG, Hackett RP, Rehder RS, Ainsworth DM, Shannon KJ, Erickson BK, Erb HN, Jansson N, Soderholm LV and Thorson LM (1997). Effects of airway obstruction on transmural pulmonary artery pressure in exercising horses. *American Journal of Veterinary Research* **58**: 897-903.
- 14 Hackett RP, Ducharme NG, Ainsworth DM, Erickson BK, Hollis NE, Soderholm LV and Thorson LM (1999). Effects of extrathoracic airway obstruction on intrathoracic pressure and pulmonary artery pressure in exercising horses. *American Journal of Veterinary Research* **60**: 485-494.
- 15 Ducharme NG, Hackett RP, Gleed RD, Ainsworth DM, Erb HN, Mitchell LM and Soderholm LV (1999). Pulmonary capillary pressure in horses undergoing alteration of pleural pressure by imposition of various upper airway resistive loads. *Equine Veterinary Journal Supplement* **30**: 27-33.
- 16 Cook WR, Williams RM, Kirker-Head CA and Verbridge DJ (1988). Upper airway obstruction (partial asphyxia) as the possible cause of exercise-induced pulmonary hemorrhage in the horse: an hypothesis. *Journal of Equine Veterinary Science* **8**: 11-26.
- 17 Poole DC, Kindig CA, Fenton G, Ferguson L, Rush BR and Erickson HH (2000). Effects of external nasal support on pulmonary gas exchange and EIPH in the horse. *Journal of Equine Veterinary Science* **20**: 579-585.
- 18 Geor RJ, Ommundson L, Fenton G and Pagan JD (2001). Effects of an external nasal strip and furosemide on pulmonary haemorrhage in Thoroughbreds following high-intensity exercise. *Equine Veterinary Journal* **33**: 577-584.
- 19 Kindig CA, McDonough P, Fenton G, Poole DC and Erickson HH (2001). Efficacy of nasal strip and furosemide in mitigating EIPH in Thoroughbred horses. *Journal of Applied Physiology* **91**: 1396-1400.
- 20 Erickson HH, Kindig CA and Poole DC (2001). Role of the airways in exercise-induced pulmonary haemorrhage. *Equine Veterinary Journal* **33**: 537-539.
- 21 Holcombe SJ, Breney C, Cornelisse CJ, Derksen FJ and Robinson NE (2002). Effect of commercially available nasal strips on airway resistance in exercising horses. *American Journal of Veterinary Research* **63**: 1101-1105.
- 22 McDonough P, Kindig CA, Hildreth TS, Padilla DJ, Behnke BJ, Erickson HH and Poole DC (2004). Effect of furosemide and the equine nasal strip on EIPH and time to fatigue in running Thoroughbred horses. *Equine and Comparative Exercise Physiology* **1**: 177-184.
- 23 Hildreth TS, McDonough P, Padilla DJ, Behnke BJ, Poole DC and Erickson HH (2003). Is immunotherapy effective in reducing exercise-induced pulmonary haemorrhage (EIPH). *The FASEB Journal* **17**: A939.
- 24 McKane SA and Slocombe RF (1999). Sequential changes in bronchoalveolar cytology after autologous blood inoculation. *Equine Veterinary Journal* **30**: 126-130.

- 25 O'Callaghan MW, Pasco JR and Tyler WS (1987). Exercise-induced pulmonary hemorrhage in the horse: results of a detailed clinical, post mortem and imaging study. V. Microscopic observations. *Equine Veterinary Journal* **19**: 411-418.
- 26 O'Callaghan MW, Pascoe JR and Tyler WS (1987). Exercise-induced pulmonary hemorrhage in the horse: results of a detailed clinical, post mortem and imaging study. VIII. Conclusions and implications. *Equine Veterinary Journal* **19**: 428-434.
- 27 McKane SA and Slocombe RF (2002). Alveolar fibrosis and changes in equine lung morphometry in response to intrapulmonary blood. *Equine Veterinary Journal Supplement* **34**: 451-458.
- 28 Newton JR and Wood JLN (2002). Evidence of an association between inflammatory airway disease and EIPH in young Thoroughbreds during training. *Equine Veterinary Journal Supplement* **34**: 417-424.
- 29 Robinson NE and Derksen FJ (1980). Small airway obstruction as a cause of exercise-associated pulmonary hemorrhage: a hypothesis. *Proceedings of the American Association of Equine Practitioners* **26**: 421-430.
- 30 Schroter RC, Marlin DJ and Denny E (1998). Exercise-induced pulmonary hemorrhage (EIPH) in horses results from locomotory impact induced trauma - a novel, unifying concept. *Equine Veterinary Journal* **30**: 186-192.
- 31 Schroter RC, Leeming A, Denny E, Bharath A and Marlin DJ (1999). Modeling impact-initiated wave transmission through lung parenchyma in relation to the aetiology of exercise-induced pulmonary haemorrhage. *Equine Veterinary Journal Supplement* **30**: 34-38.
- 32 Newton JR, Rogers K, Marlin DJ, Wood JL and Williams RB (2005). Risk factors for epistaxis on British racecourses: evidence for locomotory impact-trauma contributing to the aetiology of exercise-induced pulmonary hemorrhage. *Equine Veterinary Journal* **37**: 402-411.
- 33 Snow DH and Harris RC (1985). Thoroughbreds and Greyhounds: biochemical adaptations in creatures of nature and of man. In: Berlin-Heidelberg GR (ed.) *Circulation, Respiration and Metabolism*. Cambridge, NY: Springer-Verlag, pp. 227-239.
- 34 Nold JL, Peterson LJ and Fedde MR (1991). Physiological changes in the running greyhound (*Canis domesticus*): influence of race length. *Comparative Biochemistry and Physiology* **100A**: 623-627.
- 35 Poole DC and Erickson HH (2004). Heart and vessels: function during exercise and response to training. In: Hinchcliff KW, Kaneps AJ and Geor RJ (eds) *Equine Sports Medicine and Surgery - Basic and Clinical Sciences of the Equine Athlete*. New York: WB Saunders, pp. 699-727.
- 36 Staaden RV (1998). Section IX. Exercise and training - The exercise physiology of sporting dogs. In: Bloomberg MS, Dee JF and Taylor RA (eds) *Canine Sports Medicine and Surgery*. Philadelphia, PA: WB Saunders, pp. 357-363.
- 37 Rose RJ and Bloomberg MS (1989). Response to sprint exercise in the greyhound: effects on hematology, serum biochemistry and muscle metabolites. *Research in Veterinary Science* **47**: 212-218.
- 38 Neuhaus D, Fedde MR and Gaehtgens P (1992). Changes in haemorrheology in the racing greyhound as related to oxygen delivery. *European Journal of Applied Physiology* **65**: 278-285.
- 39 Mathieu-Costello O, Willford DC, Fu Z, Garden RM and West JB (1995). Pulmonary capillaries are more resistant to stress failure in dogs than in rabbits. *Journal of Applied Physiology* **79**: 908-917.
- 40 Birks EK, Mathieu-Costello O, Fu Z, Tyler WS and West JB (1994). Comparative aspects of the strength of pulmonary capillaries in rabbit, dog, and horse. *Respiration Physiology* **97**: 235-246.
- 41 Pieschl RL, Toll PW, Erickson BK and West JB (1992). Pulmonary artery pressure in the exercising Greyhound. *The FASEB Journal* **6**: A2048.
- 42 Ainsworth DM, Smith CA, Eicker SW, Ducharme NG, Henderson KS, Snedden K and Dempsey JA (1997). Pulmonary-locomotory interactions in exercising dogs and horses. *Respiration Physiology* **110**: 287-294.
- 43 Saibene F, Mognomi P, Aguggini G and Clement MG (1981). Work of breathing in dog during exercise. *Journal of Applied Physiology* **50**: 1087-1092.
- 44 Jayaweera AR and Ehrlich W (1988). Efficiency of work of inspiratory muscles in standing and in exercising dogs. *Chest* **94**: 1080-1085.
- 45 Kindig CA, McDonough P, Finley MR, Behnke BJ, Richardson TE, Marlin DJ, Erickson HH and Poole DC (2001). NO inhalation reduces pulmonary arterial pressure but not hemorrhage in maximally exercising horses. *Journal of Applied Physiology* **91**: 2674-2678.
- 46 Kindig CA, Ramsel C, McDonough P, Poole DC and Erickson HH (2003). Inclined running increases pulmonary haemorrhage in the Thoroughbred horse. *Equine Veterinary Journal* **35**: 581-585.
- 47 Epp TS, McDonough P, Padilla DJ, Cox JH, Poole DC and Erickson HH (2005). The effect of herbal supplementation on the severity of exercise-induced pulmonary haemorrhage. *Equine and Comparative Exercise Physiology* **2**: 17-25.
- 48 Epp TS, McDonough P, Padilla DJ, Gentile JM, Edwards KL, Erickson HH and Poole DC (2006). Exercise-induced pulmonary haemorrhage during submaximal exercise. *Equine Veterinary Journal Supplement* **37**: 502-507.
- 49 Valdez SC, Nieto JE, Spier SJ, Owens SD, Beldomenico P and Snyder JR (2004). Effect of an external nasal dilator strip on cytologic characteristics of bronchoalveolar lavage fluid in Thoroughbred racehorses. *Journal of the American Veterinary Medical Association* **224**: 558-561.
- 50 Hawkins EC and Berry CR (1999). Use of a modified stomach tube for bronchoalveolar lavage in dogs. *Journal of the American Veterinary Medical Association* **11**: 1635-1639.
- 51 Fogarty U (1990). Evaluation of a bronchoalveolar lavage technique. *Equine Veterinary Journal* **22**: 174-176.
- 52 Fogarty U and Buckley T (1991). Bronchoalveolar lavage findings in horses with exercise intolerance. *Equine Veterinary Journal* **23**: 434-437.
- 53 Mason DK, Collins EA and Watkins KL (1983). Respiratory system. In: Snow DH, Persson SGB and Rose RJ (eds) *Equine Exercise Physiology*. Cambridge, UK: Granta Editions, pp. 57-63.
- 54 Huston LJ, Bayly WM, Liggitt HD and Magnuson NS (1987). Alveolar macrophage function in Thoroughbreds after strenuous exercise. In: Gillespie JR and Robinson NE (eds) *Equine Exercise Physiology 2*. Davis, CA: ICEEP Publications, pp. 243-252.
- 55 Commins LM, Loegering DJ and Gudewicz PW (1990). Effect of phagocytosis of erythrocytes and erythrocyte ghosts on macrophage phagocytic function and hydrogen peroxide production. *Inflammation* **14**: 705-716.
- 56 Wong CW, Smith SE, Thong YH, Opdebeedk JP and Thornton JR (1992). Effects of exercise on various immune functions in horses. *American Journal of Veterinary Research* **53**: 1414-1417.
- 57 Hand WL and King-Thompson NL (1983). Effect of erythrocyte ingestion on macrophage antibacterial function. *Infection and Immunology* **40**: 917-923.
- 58 Step DL, Freeman KP, Gleed RD and Hackett RP (1991). Cytologic and endoscopic findings after intrapulmonary blood inoculation in horses. *Equine Veterinary Science* **11**: 340-345.

- 59 Raidal SL, Love DN, Bailey GD and Rose RJ (2000). The effect of high intensity exercise on the functional capacity of equine pulmonary alveolar macrophages and BAL-derived lymphocytes. *Research in Veterinary Science* **68**: 249-253.
- 60 Laegreid WW, Huston LJ, Basaraba RJ and Crisman MV (1988). The effects of stress on alveolar macrophage function in the horse: an overview. *Equine Practice* **10**: 9-16.
- 61 Custer G, Balcerzak S and Rinehart J (1982). Human macrophage hemoglobin-iron metabolism *in vitro*. *American Journal of Hematology* **13**: 23-36.
- 62 Mateos F, Brock JH and Pères-Arellano JL (1998). Iron metabolism in the lower respiratory tract. *Thorax* **53**: 594-600.
- 63 Pinsker KL, Norin AJ, Kamholz SL, Montefusco C, Schreiber K, Hagstrom JWC and Veith FJ (1980). Cell content in repetitive canine bronchoalveolar lavage. *The International Academy of Cytology Acta Cytologica* **24**: 558-583.
- 64 Rebar AH, DeNicola DB and Muggenburg BA (1980). Bronchopulmonary lavage cytology in the dog: normal findings. *Veterinary Pathology* **17**: 294-304.
- 65 Hawkins EC, DeNicola DB and Kuehn NF (1990). Bronchoalveolar lavage in the evaluation of pulmonary disease in the dog and cat. *Journal of Veterinary Internal Medicine* **4**: 267-274.
- 66 Mayer P, Laber G and Walzl H (1990). Bronchoalveolar lavage in dogs; analysis of proteins and respiratory cells. *Journal of Veterinary Medicine* **37**: 392-399.
- 67 Vail DM, Mahler PA and Soergel SA (1995). Differential cell analysis and phenotypic subtyping of lymphocytes in bronchoalveolar lavage fluid from clinically normal dogs. *American Journal of Veterinary Research* **56**: 282-285.
- 68 McCauley M, Atwell RB, Sutton RH and Lumsden JS (1998). Unguided bronchoalveolar lavage techniques and residual effects in dogs. *Australian Veterinary Journal* **76**: 161-165.
- 69 Rha JY and Mahony O (1999). Bronchoscopy in small animal medicine: indications, instrumentation, and techniques. *Clinical Techniques in Small Animal Practice* **14**: 207-212.
- 70 McCullough S and Brinson J (1999). Collection and interpretation of respiratory cytology. *Clinical Techniques in Small Animal Practice* **14**: 220-226.
- 71 Rajamaki MM, Jarvinen AK, Saari SAM and Maisi PS (2001). Effect of repetitive bronchoalveolar lavage on cytologic findings in healthy dogs. *American Journal of Veterinary Research* **62**: 13-16.
- 72 Andreasen CB (2003). Bronchoalveolar lavage. In: Cowell RL (ed.) *The Veterinary Clinics of North America - Small Animal Practice Cytology II*. vol. **33**. New York: WB Saunders, pp. 69-88.
- 73 Kwak SH, Choi JI and Park JT (2004). Effects of propofol on endotoxin-induced acute lung injury in rabbit. *Journal of Korean Medical Science* **19**: 55-61.
- 74 Takao Y, Mikawa K, Nishina K and Obara H (2005). Attenuation of acute lung injury with propofol in endotoxemia. *Anesthesia and Analgesia* **100**: 810-816.
- 75 Davis MS, McKiernan B, McCullough S, Nelson S and Mandstagr RE (2002). Racing Alaskan sled dogs as a model of "Ski Asthma". *American Journal of Respiratory Critical Care Medicine* **166**: 878-882.
- 76 Olsen SC, Coyne CP, Lowe BS, Pelletier N, Raub EM and Erickson HH (1992). Influence of furosemide on hemodynamic responses during exercise in horses. *American Journal of Veterinary Research* **53**: 742-747.
- 77 Fedde MR and Erickson HH (1998). Increase in blood viscosity in the sprinting horse: can it account for the high pulmonary arterial pressure? *Equine Veterinary Journal* **30**: 329-334.
- 78 Schneider HP, Truex RC and Knowles JO (1964). Comparative observations of the hearts of Mongrel and Greyhound dogs. *Anatomical Record* **149**: 173-180.
- 79 Steel JD, Taylor RI, Davis PE, Stewart GA and Salmon PW (1976). Relationships between heart score, heart weight, and body weight in greyhound dogs. *Australian Veterinary Journal* **52**: 561-564.
- 80 Staaden R (1980). Cardiovascular System of the racing dog. In: Kirk RW (ed.) *Current Veterinary Therapy VII, Small Animal Practice*. Philadelphia, PA: WB Saunders Company, pp. 349.
- 81 Schoning P, Erickson HH and Milliken GA (1995). Body weight, heart weight, and heart-to-body weight ratio in Greyhounds. *American Journal of Veterinary Research* **56**: 420-422.
- 82 Rose RJ, Hodgson DR, Kelso TB, McCutcheon LJ, Reid TA, Bayly WM and Gollnick PD (1988). Maximum O₂ uptake, O₂ debt and deficit, and muscle metabolites in Thoroughbred horses. *Journal of Applied Physiology* **64**: 781-788.
- 83 Evans DL and Rose RJ (1988). Determination and repeatability of maximum oxygen uptake and other cardiorespiratory measurements in the exercising horse. *Equine Veterinary Journal* **20**: 94-98.
- 84 Birks EK, Mathieu-Costello O, Fu Z, Tyler WS and West JB (1997). Very high pressures are required to cause stress failure of pulmonary capillaries in Thoroughbred racehorses. *Journal of Applied Physiology* **82**: 1584-1592.
- 85 Weibel ER, Marques LB, Constantinopol M, Duffey F, Gehr P and Taylor CR (1987). Adaptive variation in the mammalian respiratory system in relation to energetic demand: VI. The pulmonary gas exchanger. *Respiration Physiology* **69**: 81-100.
- 86 Marlin DJ, Schroter RC, Cashman PMM, Deaton CM, Poole DC, Kindig CA, McDonough P and Erickson HH (2002). Movements of thoracic and abdominal compartments during ventilation at rest and during exercise. *Equine Veterinary Journal Supplement* **34**: 383-390.
- 87 Elliot AR, Fu Z, Tsukimoto K, Prediletto R, Mathieu-Costello O and West JB (1992). Short-term reversibility of ultrastructural changes in pulmonary capillaries caused by stress failure. *Journal of Applied Physiology* **73**: 1150-1158.
- 88 Hillidge CJ, Lane TJ, Johnson EL and Asquith RL (1984). Preliminary investigations of exercise-induced pulmonary hemorrhage in racing Quarter Horses. *Journal of Equine Veterinary Science* **4**: 21-23.
- 89 Ghio AJ, Ghio C and Bassett M (2006). Exercise-induced pulmonary hemorrhage after running a marathon. *Lung* **184**: 331-333.
- 90 Newton JR, Rogers K, Marlin DJ, Wood JLN and Williams RB (2005). Risk factors for epistaxis on British racecourses: evidence for locomotory impact-induced trauma contributing to the aetiology of exercise-induced pulmonary haemorrhage. *Equine Veterinary Journal* **37**: 402-411.
- 91 Hinchcliff KW (2000). Counting red cells - is it the answer to EIPH? *Equine Veterinary Journal* **32**: 362-363.
- 92 Sweeney CR, Rossier Y, Ziemer EL and Lindborg S (1992). Effects of lung site and fluid volume on results of bronchoalveolar lavage fluid analysis in horses. *American Journal of Veterinary Research* **53**: 1376-1379.
- 93 Robinson NE, Derksen FJ, Slocombe RF and Scott JS (1986). Bronchoalveolar lavage. In: Huntingdon P (ed.) *Proceedings for the 8th Bain-Fallon Memorial Lectures*. Australia: AEVA, pp. 6-12.
- 94 McKane SA and Rose RJ (1993). Radiographic determination of the location of a blindly placed bronchoalveolar lavage catheter. *Equine Veterinary Journal* **5**: 329-332.

- 95 McKane SA and Rose RJ (1995). Effects of exercise intensity and training on bronchoalveolar lavage cytology. *Equine Veterinary Journal Supplement* **18**: 58-62.
- 96 Carre PH, Laviolette M, Belanger J and Cormier Y (1985). Technical variations of bronchoalveolar lavage (BAL): influence of atelectasis and the lung region lavaged. *Lung* **163**: 117-125.
- 97 Whaley SL, Muggenburg BA, Seiler FA and Wolff RK (1987). Effect of aging on tracheal mucociliary clearance in beagle dogs. *Journal of Applied Physiology* **62**: 1331-1334.
- 98 Willoughby RA, Ecker GL, McKee SL and Riddolls LJ (1991). Use of scintigraphy for the determination of mucociliary clearance rates in normal, sedated, diseased, and exercised horses. *Canadian Journal of Veterinary Research* **55**: 315-320.
- 99 Clark CK, Lester GD, Vetro T and Rice B (1995). Bronchoalveolar lavage in horses: effect of exercise and repeated sampling on cytology. *Australian Veterinary Journal* **72**: 249-252.
- 100 Kindig CA, Gallatin LL, Erickson HH, Fedde MR and Poole DC (2000). Cardiorespiratory impact of the nitric oxide synthase inhibitor L-NAME in the exercising horse. *Respiration Physiology* **120**: 151-166.
- 101 Boothe HW, Booth DM, Komkov A, Longnecker MT and Hightower D (1993). Tracheal mucociliary transport rate in awake dogs. *American Journal of Veterinary Research* **54**: 1812-1816.
- 102 Couëtil LL and Hinchcliff KW (2004). Chapter 29. Non-infectious respiratory diseases of the lower respiratory tract. In: Hinchcliff KW, Kaneps AJ and Geor RJ (eds) *Equine Sports Medicine and Surgery - Basic and Clinical Sciences of the Equine Athlete*. New York: WB Saunders, pp. 613-656.

As well, the harder the exercise, the greater the increase in pulmonary capillary pressure and the probability of blood vessel rupture. Studies conducted by researchers at the University of California, Davis, and the Japan Racing Association have shed some light on why blood pressures in the lung increase so dramatically during exercise. To understand their findings, we first need a brief review of the flow of blood through the heart and lungs. Hinchcliff, K.W. Effects of furosemide on athletic performance and exercise-induced pulmonary hemorrhage in horses. *Journal of the American Veterinary Medical Association*, 215, 630-635, 1999. Of 235 Thoroughbred racehorses examined with a flexible fiberoptic endoscope within 2 hours of racing to determine the frequency of exercise-induced pulmonary hemorrhage (EIPH), 103 (43.8%) had various degrees of hemorrhage in the tracheal lumen. Two of these horses (0.8%) subsequently had blood flow from the nostrils. Blood seemed to originate from the lung. Statistical analysis of frequency data for 191 horses which finished in 1st, 2nd, and 3rd places did not show any relationship between EIPH and horse's age, sex, or finishing position. However, a trend toward an increased frequency of Post-mortem examination of 74 racing greyhounds with sudden, unexpected death was undertaken at the Veterinary School, University of Nottingham, UK, from 2015 to 2019. Despite the acute clinical event, three cases demonstrated histological evidence of previous muscle damage with muscular regeneration, fibrosis and evidence of chronic haemorrhage. To our knowledge, this is the first detailed description of exercise-induced trauma to the iliopsoas muscle leading to extensive haemoretroperitoneum and haemoabdomen with subsequent hypovolaemic shock and death in racing greyhounds. Download full-text PDF. Source. racing's very core. That exercise-induced pulmonary hemorrhage (EIPH) occurs is a medical, and physiological reality. That every reasonable exigency is not taken to reduce/prevent it would. Figure 7 Left: Schematic showing exercise-induced pulmonary hemorrhage in the horse. Right: Scanning electron micrograph of the cranial lobe of an exercised pony reveals red blood cells (R) and proteinaceous material (P) in the alveoli. Exercise-induced pulmonary hemorrhage. High pulmonary capillary intravascular pressures. Exercise-induced pulmonary hemorrhage: where are we now? Investigations using BAL indicate that hemorrhage occurs in almost all horses in racing or training, 2,60,61 and BAL cytological findings cohere closely with clinical disease, pulmonary histopathology and endoscopic evidence of EIPH. 57-60 Concerns regarding BAL most often include the need for rapid access to the horse post-race, the need for sedation (eg, 10-20 µg/kg intravenous detomidine).